
Physics with neutrons 2

Michael Leitner, michael.leitner@frm2.tum.de

Summer semester 2017

Exercise sheet 5

To be discussed 2017-05-30, room C.3203

Franz Haslbeck, franz.haslbeck@frm2.tum.de

EXERCISE 5.1

The Debye–Waller factor is used in to describe the attenuation of coherent neutron scattering caused by thermal motion:

$$f_{\text{DWF}} = e^{-Q^2/3\langle u^2 \rangle} = e^{-2W(\mathbf{Q})}.$$

For a cubic Bravais lattice we can make the following approximation:

$$2W = \frac{Q^2 \hbar}{6MN} \int \frac{Z(\omega)}{\omega} \coth\left(\frac{\hbar\omega}{2k_B T}\right) d\omega,$$

where $Z(\omega)$ is the phonon density of states, M is the mass of the atom and N is the number of atoms in the crystal.

Within the Debye approximation, when the velocity of sound is frequency independent, we can express the phonon density of states for a cubic crystal with side length L by (in analogy with the theory of the black body radiation):

$$Z(\omega) = \frac{1}{2\pi^2} L^3 \left(\frac{1}{c_L^3} + \frac{2}{c_T^3} \right) \omega^2. \quad (1)$$

c_L and c_T are the longitudinal and transverse velocity of sound, respectively. The total number of normal modes is $3N$. Therefore, we can put:

$$3N = \int_0^{\Omega_D} Z(\omega) d\omega. \quad (2)$$

ω_D is the maximum frequency of the normal mode and $\omega_D = \frac{\hbar\omega_D}{k_B}$ is the Debye temperature.

1. Calculate ω_D from the equations (1) and (2).
2. Calculate the asymptotic behaviour of $2W$ for $T \ll \Theta_D$ and $T \gg \Theta_D$.
3. Copper crystallizes in fcc-lattice ($a = 3.615 \text{ \AA}$, $\rho_{\text{Cu}} = 8920 \text{ kg/m}^3$, $c_L = 4760 \text{ m/s}$ and $c_T = 2320 \text{ m/s}$);
 - a) Calculate θ_D and show that the obtained value is reasonable.
 - b) Figure 1 and 2 show $Z(\omega)$ and the dispersion relation for copper, respectively (Nilsson 1973). What are the reasons of deviations to the Debye model?




Figure 1: Phonon frequency distributions calculated from the Born-von Kármán models.




Figure 2: Dispersion curves for Cu at 80K.

4. Calculate the mean amplitude $\langle u^2 \rangle$ for copper at $T = 20K, 100K, 500K, 1000K$.
5. Estimate the attenuation in a neutron powder diffraction measurement with wavelength $\lambda = 1.188$ Å of (100) and (440) reflex due to the Debye-Waller factor ($T = 200K$).
6. a) Why do soft materials have a larger Debye-Waller factor than condensed matter?
b) What is the influence of mass on the Debye-Waller factor?